



## MEETING ABSTRACT

## Open Access

# Comparing smoking habits and tobacco-related education between Canadian and Greek medical students

Georgios-Marios Pantsidis<sup>1\*</sup>, Dimitra-Iosifina Papageorgiou<sup>2</sup>, Demosthenes Bouros<sup>1</sup>

From 11th Annual Conference of the International Society for the Prevention of Tobacco Induced Diseases (ISPTID)

Athens, Greece. 9-11 December 2013

## Background

According to a survey on Canadian medical students' smoking habits and beliefs, the key results show that the prevalence of smoking among the future healthcare professionals is high and they lack of tobacco-related education [1]. Last year a similar survey was conducted at Democritus University of Thrace [2]. Its findings show that there is difference in smoking habits between the two students' groups, but their tobacco-related education is equally poor.

## Materials and methods

In both researches participated undergraduate students who completed a questionnaire about their smoking habits, attitudes and education level towards tobacco cessation interventions.

## Results

The prevalence of cigarette smoking among Greek medical students is higher than the Canadians (24% vs. 3.3%). Although Canadian students smoke, also, other tobacco products (cigars, water pipe), the total prevalence is 15.3%. 65.5% of the Greek medical students report that they had ever tried cigarettes, but only 29.9% of the Canadian students make a same statement. Both students groups reported that they have moderate levels of education concerning tobacco-related subjects and cessation techniques. Only 8.1% of Greek and 10% of Canadian medical students report that they had ever received trainings in smoking cessation methods. Finally

only a small percentage seems to be familiar with the cessation guidelines and only a few students are aware of the fact that they lack knowledge to help their patients cease smoking.

## Conclusions

The prevalence of cigarettes smoking among Greek medical students is significantly higher. Also the tobacco-related education in both countries is equally poor. It is desperately necessary to enhance the medical schools' curricula with courses regarding smoking issues, since future physicians have a key-role in tobacco cessation and prevention.

## Authors' details

<sup>1</sup>Faculty of Medicine, Democritus University of Thrace, Alexandroupolis, 68100, Greece. <sup>2</sup>Faculty of Medicine, University of Thessaly, Larisa, 41110, Greece.

Published: 6 June 2014

## References

1. Vanderhoek AJ, Hammal F, Chappell A, Wild TC, Raupach T, Finegan BA: Future physicians and tobacco: An online survey of the habits, beliefs and knowledge base of medical students at a Canadian university. *Tobacco Induced Diseases* 2013, **11**(1):9.
2. Pantsidis GM, Papageorgiou DI, Bouros D: Smoking habits, attitudes and training among medical students of the Democritus University of Thrace. *Pneumon* 2012, **25**:208-218.

doi:10.1186/1617-9625-12-S1-A5

**Cite this article as:** Pantsidis et al.: Comparing smoking habits and tobacco-related education between Canadian and Greek medical students. *Tobacco Induced Diseases* 2014 **12**(Suppl 1):A5.

\* Correspondence: [george.marios.pantsidis@gmail.com](mailto:george.marios.pantsidis@gmail.com)<sup>1</sup>Faculty of Medicine, Democritus University of Thrace, Alexandroupolis, 68100, Greece

Full list of author information is available at the end of the article